



GREENVILLE
CENTRAL SCHOOL DISTRICT

PO Box 129
Greenville, NY 12083
(518) 966-5070 ext. 305
Fax: (518) 966-6033

HS/MS Nurse 518-966-5070 Ext. 420
Fax: 518-966-5338

Scott M. Ellis Nurse 518-966-5070 Ext. 320
Fax: 518-966-5321

HEALTH CERTIFICATE / APPRAISAL FORM

For All New Entrants and Students Entering Grades UPK, K, 2, 4, 7, and 10

Completed by Physician

Name: _____ **Date of Exam:** _____
Date of Birth: _____
School: _____ **Gender:** M F **Grade:** _____

IMMUNIZATIONS / HEALTH HISTORY

- Immunization record attached
 No immunizations given today
 Immunizations given since last Health Appraisal:
- Sickle Cell Screen: Positive Negative Not done Date: _
PPD: Positive Negative Not done Date: _
Elevated Lead: Yes No Not done Date: _
Dental Referral Yes No Not done Date: _

Significant Medical/Surgical History: See attached _____

Specify current diseases: Asthma** Diabetes**
 Type 1 Type 2
 Hyperlipidemia Hypertension
 Other: _____

**** List All Medications/Treatments on page 2 of Health Appraisal Form****

Allergies: LIFE THREATENING Food: _____ Insect: _____ Other: _____
 Seasonal Medication: _____

PHYSICAL EXAM

Height: _____ Weight: _____ Blood Pressure: _____

REQUIRED:

REQUIRED:

Referral

Body Mass Index: _____ . _____	Vision - without glasses/contact lenses	R	L	
Weight Status Category (BMI Percentile):	Vision - with glasses/contact lenses	R	L	
<input type="checkbox"/> less than 5 th <input type="checkbox"/> 5 th through 49 th <input type="checkbox"/> 50 th through 84 th	Vision - Near Point	R	L	
<input type="checkbox"/> 85 th through 94 th <input type="checkbox"/> 95 th through 98 th <input type="checkbox"/> 99 th and higher	Hearing <input type="checkbox"/> Pass 20 db sc both ears or:	R	L	

EXAM ENTIRELY NORMAL Tanner: I. II. III. IV. V. Scoliosis: Negative Positive: _____

Specify any abnormality (use reverse of form if needed): _____

This exam complies with NYSED requirements above and is valid for twelve months, with the exception of any illness or injury lasting more than five days that will require review by private healthcare provider and the school medical director. Rev. 11/2/09

MEDICATIONS

Medications (list all): None Additional medications listed on reverse of form

Name: _____ Dosage/Time: _____

Name: _____ Dosage/Time: _____

Name: _____ Dosage/Time: _____

If AM dose is missed at home:

I assess this student to be self-directed Yes No

Student may self carry and self administer medication Yes No

Note: Nurse will also assess self-direction for the school setting. Please advise parent to send in additional medication in the event that emergency sheltering is necessary at school or if the morning medication has not been given.

PHYSICAL EDUCATION / SPORTS / PLAYGROUND / WORK QUALIFICATION / CSE CONSIDERATION

Physically qualified for all physical education, sports, playground, work & school activities OR only as checked:

___ Limited contact: cheerleading, gymnastics, ski, volleyball, cross-country, handball, fence, baseball, floor hockey, softball.

___ Non-contact: badminton, bowl, golf, swim, table tennis, tennis, archery, weight train, crew, dance, track, run, walk, rope jump.

Specify medical accommodations needed for school: _____ None

Known or suspected disability: _____ Please monitor

Restrictions: _____ Please monitor

Protective equipment required: Athletic Cup Sport goggles/impact resistant eyewear
 Other: _____

Physician's Signature: _____ Phone: _____ (Stamp Below)

Physician's Name/Address: _____ Fax: _____

Notes: _____

2016-17 School Year New York State Immunization Requirements for School Entrance/Attendance¹

NOTES:
Children in a prekindergarten setting should be age-appropriately immunized. The number of doses depends on the schedule recommended by the Advisory Committee on Immunization Practices (ACIP). For grades Pre-k through 8, intervals between doses of vaccine should be in accordance with the ACIP-recommended immunization schedule for persons 0 through 18 years of age. (Exception: intervals between doses of polio vaccine need to be reviewed only for grades prekindergarten, kindergarten, 1, 2, 6, 7 and 8.) Doses received before the minimum age or intervals are not valid and do not count toward the number of doses listed below. Intervals between doses of vaccine DO NOT need to be reviewed for grades 9 through 12. See footnotes for specific information for **each** vaccine. Children who are enrolling in grade-less classes should meet the immunization requirements of the grades for which they are age equivalent.

Dose requirements MUST be read with the footnotes of this schedule.

Vaccines	Prekindergarten (Day Care, Head Start, Nursery or Pre-k)	Kindergarten and Grades 1 and 2	Grades 3, 4 and 5	Grades 6, 7 and 8	Grades 9, 10, 11 and 12
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis vaccine (DTaP/DTP/Tdap)²	4 doses	5 doses or 4 doses if the 4th dose was received at 4 years of age or older or 3 doses if aged 7 years or older and the series was started at 1 year of age or older		3 doses	
Tetanus and Diphtheria toxoid-containing vaccine and Pertussis vaccine booster (Tdap)³	Not applicable			1 dose	
Polio vaccine (IPV/OPV)⁴	3 doses	4 doses or 3 doses if the 3rd dose was received at 4 years of age or older	3 doses	4 doses or 3 doses if the 3rd dose was received at 4 years of age or older	3 doses
Measles, Mumps and Rubella vaccine (MMR)⁵	1 dose	2 doses			
Hepatitis B vaccine⁶	3 doses	3 doses or 2 doses of adult hepatitis B vaccine (Recombivax) for children who received the doses at least 4 months apart between the ages of 11 through 15 years of age			
Varicella (Chickenpox) vaccine⁷	1 dose	2 doses	1 dose	2 doses	1 dose
Meningococcal conjugate vaccine (MenACWY)⁸	Not applicable			By Grade 7: 1 dose	Grade 12: 2 doses or 1 dose if the dose was received at 16 years of age or older
Haemophilus influenzae type b conjugate vaccine (Hib)⁹	1 to 4 doses	Not applicable			
Pneumococcal Conjugate vaccine (PCV)¹⁰	1 to 4 doses	Not applicable			

1. Demonstrated serologic evidence of measles, mumps, rubella, hepatitis B, varicella or polio (for all three serotypes) antibodies is acceptable proof of immunity to these diseases. Diagnosis by a physician, physician assistant or nurse practitioner that a child has had varicella disease is acceptable proof of immunity to varicella.
2. Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine. (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive a 5-dose series of DTaP vaccine at ages 2 months, 4 months, 6 months and at 15 through 18 months and at 4 years of age or older. The fourth dose may be received as early as age 12 months, provided at least 6 months have elapsed since the third dose. However, the fourth dose of DTaP need not be repeated if it was administered at least 4 months after the third dose of DTaP. The final dose in the series must be received on or after the fourth birthday.
 - b. If the fourth dose of DTaP was administered at age 4 years or older, the fifth (booster) dose of DTaP vaccine is not required.
 - c. For children born before 1/1/2005, only immunity to diphtheria is required and doses of DT and Td can meet this requirement.
 - d. Children ages 7 through 10 years who are not fully immunized with the childhood DTaP vaccine series should receive Tdap vaccine as the first dose in the catch-up series; if additional doses are needed, use Td vaccine. A Tdap vaccine (or incorrectly administered DTaP vaccine) received at 7 years or age or older will meet the 6th grade Tdap requirement.
 - e. For children 7 years of age or older who received the first dose on or after their first birthday, the immunization requirement is 3 doses. If the first dose was received before their first birthday, then 4 doses are required.
3. Tetanus and diphtheria toxoids and acellular pertussis (Tdap) vaccine. (Minimum age: 7 years)
 - a. Students 11 years of age or older entering grades 6 through 12 are required to have one dose of Tdap. A dose received at 7 years of age or older will meet this requirement.
 - b. Students who are 10 years old in grade 6 and who have not yet received a Tdap vaccine are in compliance until they turn 11 years of age.
4. Poliovirus vaccine (IPV/OPV). (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive a series of IPV at ages 2 months, 4 months and at 6 through 18 months, and 4 years of age or older. The final dose in the series must be received on or after the fourth birthday and at least 6 months after the previous dose.
 - b. For students who received their fourth dose before age 4 and prior to August 7, 2010, 4 doses separated by at least 4 weeks is sufficient.
 - c. If the third dose of polio vaccine was received at age 4 years or older and at least 6 months after the previous dose, the fourth dose of polio vaccine is not required.
5. Measles, mumps, and rubella (MMR) vaccine. (Minimum age: 12 months)
 - a. The first dose of MMR vaccine must have been received on or after the first birthday. The second dose must have been received at least 28 days (4 weeks) after the first dose to be considered valid.
 - b. Students in grades kindergarten through 12 must have received 2 doses of measles-containing vaccine, 2 doses of mumps-containing vaccine and at least 1 dose of rubella-containing vaccine.
 - c. One dose of MMR is required for prekindergarten.
6. Hepatitis B vaccine
 - a. Dose 1 may be given at birth or anytime thereafter. Dose 2 must be given at least 4 weeks (28 days) after dose 1. Dose 3 must be at least 8 weeks after dose 2 AND at least 16 weeks after dose 1 AND no earlier than 24 weeks of age.
 - b. Two doses of adult hepatitis B vaccine (Recombivax) received at least 4 months apart at age 11 through 15 years will meet the requirement.
7. Varicella (chickenpox) vaccine. (Minimum age: 12 months)
 - a. The first dose of varicella vaccine must have been received on or after the first birthday. The second dose must have been received at least 28 days (4 weeks) after the first dose to be considered valid.
 - b. For children aged less than 13 years, the recommended minimum interval between doses is 3 months (if the second dose was administered at least 4 weeks after the first dose, it can be accepted as valid); for persons aged 13 years and older, the minimum interval between doses is 4 weeks.
8. Meningococcal conjugate vaccine (MenACWY). (Minimum age: 6 weeks)
 - a. One dose of meningococcal conjugate vaccine (Menactra or Menevo) is required for students entering grade 7.
 - b. For students in grade 12, if the first dose of meningococcal conjugate vaccine was received at age 16 years or older, the second (booster) dose is not required.
9. Haemophilus influenzae type b (Hib) conjugate vaccine. (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive Hib vaccine at 2 months, 4 months, 6 months and at 12 through 15 months. Children older than 15 months must get caught up according to the ACIP catch-up schedule. The final dose must be received on or after 12 months of age.
 - b. If 2 doses of vaccine were received before 12 months of age, only 3 doses are required with dose 3 at 12 through 15 months of age and at least 8 weeks after dose 2.
 - c. If dose 1 was received at ages 12 through 14 months of age, only 2 doses are required with dose 2 at least 8 weeks after dose 1.
 - d. If dose 1 was received at 15 months of age or older, only 1 dose is required.
 - e. Hib vaccine is not required for children 5 years of age or older.
10. Pneumococcal conjugate vaccine (PCV). (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive PCV vaccine at 2 months, 4 months, 6 months and at 12 through 15 months. Children older than 15 months must get caught up according to the ACIP catch-up schedule. The final dose must be received on or after 12 months of age.
 - b. Unvaccinated children 7 through 11 months of age are required to receive 2 doses, at least 4 weeks apart, followed by a third dose at age 12 through 15 months.
 - c. Unvaccinated children 12 through 23 months of age are required to receive 2 doses of vaccine at least 8 weeks apart.
 - d. If one dose of vaccine was received at 24 months of age or older, no further doses are required.
 - e. For further information, refer to the PCV chart available in the School Survey Instruction Booklet at: www.health.ny.gov/prevention/immunization/schools

For further information contact:

**New York State Department of Health
Bureau of Immunization
Room 649, Corning Tower ESP
Albany, NY 12237
(518) 473-4437**

**New York City Department of Health and Mental Hygiene
Program Support Unit, Bureau of Immunization,
42-09 28th Street, 5th floor
Long Island City, NY 11101
(347) 396-2433**