

Vaccines*	Grade K	Grades 1-5	Grade 6	Grade 7-12
Polio vaccine (IPV/OPV)	3-4 doses <sup>1</sup>	3 doses	3-4 doses <sup>2</sup>	3 doses
Tdap (Tetanus & Diphtheria Toxoid-containing vaccine and Pertussis vaccine booster)	Not required	Not required	1 dose <sup>3</sup>	1 dose <sup>3</sup>
DTap/DTP	4-5 doses <sup>4</sup>	4-5 doses <sup>4</sup>	3 doses	3 doses
HEP B (Hepatitis B Vaccine <sup>6</sup> )	3 doses	3 doses	3 doses	3 doses
MMR (Measles, Mumps and Rubella)	1 dose	2 doses	2 doses	2 doses
Varicella (Chickenpox)	2 doses	1 dose	2 doses	1 dose

<sup>1</sup> If 4 or more doses administered before age 4, additional dose should be received at age 4 to 6

<sup>2</sup> If OPV & IPV were administered as part of a series, a total of 4 doses should be received, regardless of child's current age

<sup>3</sup> If child received Tdap vaccine at age 7 or older, this booster is not required

<sup>4</sup> If 4<sup>th</sup> dose of vaccine was administered at age 4 or older, the fifth dose is not necessary

\*Demonstrated serologic evidence of measles, mumps, rubella, hepatitis B, varicella or Polio (for all three serotypes) antibodies is acceptable proof of immunity to these diseases. Diagnosis by a physician, physician assistant or nurse practitioner that a child has had varicella disease is acceptable proof of immunity to varicella.